



# A new modified King–Werner method for solving nonlinear equations<sup>☆</sup>

Liang Chen<sup>a,\*</sup>, Yanfang Ma<sup>b</sup>

<sup>a</sup> School of Mathematical Science, Huaibei Normal University, Huaibei 235000, Anhui, PR China

<sup>b</sup> School of Computer Science and Technology, Huaibei Normal University, Huaibei 235000, PR China

## ARTICLE INFO

### Article history:

Received 9 December 2010

Received in revised form 21 August 2011

Accepted 6 September 2011

### Keywords:

Nonlinear equations

Secant method

Iterative method

Root-finding

King–Werner method

## ABSTRACT

In this paper, a new method for solving nonlinear equations  $f(x) = 0$  is presented. Analysis of the convergence shows that the asymptotic convergence order of this method is  $1 + \sqrt{3}$ . Some numerical results are given to demonstrate its efficiency.

Crown Copyright © 2011 Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

Numerical methods for solving nonlinear equations are a popular and important research topic in numerical analysis. In this paper, we consider iterative methods to find a simple root of a nonlinear equation  $f(x) = 0$ , where  $f : D \subset \mathbb{R} \rightarrow \mathbb{R}$ , for an open interval  $D$ , is a scalar function.

We all know that Newton's method is an important and basic approach for solving nonlinear equations [1–5], and its formulation is given by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}. \quad (1)$$

This method is a quadratic method.

To improve the local order of convergence, a number of modified methods have been studied and reported in the literature [6–14]. By employing a second derivative evaluation we can obtain some well-known third-order methods, such as Chebyshev's method, Halley's method [6] and the super-Halley method [8]. However, in many other cases, it is expensive to compute the derivative, and the above methods are still restricted in practical applications. The well known secant method is given by

$$x_{n+1} = x_n - \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} f(x_n). \quad (2)$$

The method does not require any derivative, but its order is only 1.618. To improve this method, many modified methods called the secant-like method have been proposed in [9,15,16].

<sup>☆</sup> The work is supported by Shanghai Natural Science Foundation (NO. 10ZR1410900), Key Disciplines of Shanghai Municipality (NO. S30104), Natural Science Research Funds of Anhui Provincial for Universities (Grant No. KJ2011A248) and the Open Fund of Shanghai Key Laboratory of Trustworthy Computing.

\* Corresponding author. Tel.: +86 13774415529.

E-mail addresses: [clmyf2@163.com](mailto:clmyf2@163.com), [chenliang1977@gmail.com](mailto:chenliang1977@gmail.com) (L. Chen).

Recently, Zhang et al. [15] and Wang et al. [16] have proposed a new two-step secant-like method and a new three-step secant-like method respectively, and the convergence order of these methods are improved. Zhang's method proposed in [15] is very similar to the well-known King–Werner method with order 2.414 [17], which has the form [18,19]:

$$\begin{cases} x_{n+1} = x_n - f' \left( \frac{x_n + y_n}{2} \right)^{-1} f(x_n), \\ y_{n+1} = x_{n+1} - f' \left( \frac{x_n + y_n}{2} \right)^{-1} f(x_{n+1}). \end{cases} \quad (3)$$

Here, we attempt to improve the order of the classical method defined by (1), and (2) by using previous information, and then we present a new iterative method for solving nonlinear equations. Analysis of the convergence shows that the asymptotic convergence order of this method is  $1 + \sqrt{3}$  which is equal to the method defined by Wang et al. proposed in [16]. Per iteration, the methods require two evaluations of the function and one of its first derivative and therefore the efficiency, in term of function evaluations, of the new methods is equal to  $\sqrt[3]{1.618} \approx 1.17398$ , which is better than the method proposed in [16]. Finally, some numerical examples are also given.

## 2. Notation and basic results

Let  $f(x)$  be a real function with a simple root  $x^*$  and let  $\{x_n\}_{n=0}^{\infty}$  be a sequence of real numbers that converges to  $x^*$ . We say that the *order of convergence* is  $q$  if there exists a  $q \in \mathbb{R}^+$  such that

$$\lim_{n \rightarrow +\infty} \frac{|x_{n+1} - x^*|}{|x_n - x^*|^q} = C \neq 0, \infty.$$

Let  $e_n = x_n - x^*$  be the  $n$ -th iterate error. We call

$$e_{n+1} = Ce_n^q + \dots, \quad (4)$$

the *error equation*. If we can obtain the error equation for the method, then the value of  $q$  is its order of convergence.

## 3. The new method

Here, in order to construct our method, we use the following second-order polynomial function proposed in [16]:

$$P(x) = f(x_n) + v_n^{-1}(x - x_n) + \frac{(v_{n-1}^{-1} - v_n^{-1})(x - x_n)(x - y_n)}{\alpha_1 x_{n-1} + \alpha_2 y_{n-1} + (2 - \alpha_1 - \alpha_2)z_{n-1} - \beta_1 x_n - \beta_2 y_n - (2 - \beta_1 - \beta_2)z_n} \quad (5)$$

where  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{R}$ , and  $y_n = x_n - v_{n-1}f(x_n)$ ,  $v_n = f' \left( \frac{x_n + y_n}{2} \right)^{-1}$  and  $z_n = x_n - v_n f(x_n)$ , respectively.

It is easy to obtain that

$$(v_{n-1}^{-1} - v_n^{-1})(y_n - x_n)(z_n - y_n) = v_n^{-1}(y_n - z_n)^2.$$

In order to eliminate the nonlinearity, we replace  $x_{n+1}$  in the terms  $(x_{n+1} - x_n)$  and  $(x_{n+1} - y_n)$  of  $P(x_{n+1})$  with  $y_n$  and  $z_n$ , respectively. By solving the function (5), we can obtain a new method

$$\begin{cases} y_n = x_n - v_{n-1}f(x_n), \\ v_n = f' \left( \frac{x_n + y_n}{2} \right)^{-1} \\ z_n = x_n - v_n f(x_n), \\ x_{n+1} = z_n - \frac{(y_n - z_n)^2}{\alpha_1 x_{n-1} + \alpha_2 y_{n-1} + (2 - \alpha_1 - \alpha_2)z_{n-1} - \beta_1 x_n - \beta_2 y_n - (2 - \beta_1 - \beta_2)z_n} \end{cases} \quad (6)$$

where  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{R}$ .

The method defined by (6) can be viewed as an iterative method with three substeps. The first two substeps are the well-known King–Werner method [18,19]. The third substep is an acceleration by using the values computed previously.

At the beginning of the process, the values of  $x_0, y_0$  and  $v_{-1}$  need to be given by some approaches. The choice of  $v_{-1}$  cannot affect the asymptotic convergence order of the method defined by (6) while  $v_{-1} \neq 0$ . Since the first iteration cannot carry out the third substep, and hence we let  $x_1 = z_0$ .

**Theorem 1.** Assume that the function  $f : D \subset \mathbb{R} \rightarrow \mathbb{R}$  for an open interval  $D$  has a simple root  $x^* \in D$ . Let  $f(x)$  have first, second and third derivatives in the interval  $D$ , then the asymptotic convergence of the method defined by (6) is  $1 + \sqrt{3}$  when  $\alpha_1 = \alpha_2 = 1$ .

**Proof.** Let  $d_n = y_n - x^*$  and  $w_n = z_n - x^*$ . Using Taylor expansion, we get

$$f' \left( \frac{x_n + y_n}{2} \right) = f'(x^*) \left[ 1 + C_2(e_n + d_n) + \frac{3}{4}C_3(e_n + d_n)^2 + \frac{1}{2}C_4(e_n + d_n)^3 \cdots \right] \quad (7)$$

where  $C_k = \frac{1}{k!} \frac{f^{(k)}(x^*)}{f'(x^*)}$ ,  $k = 2, 3, \dots$  we have

$$\begin{aligned} v_n &= f' \left( \frac{x_n + y_n}{2} \right)^{-1} \\ &= \frac{1}{f'(x^*)} \frac{1}{1 + C_2(e_n + d_n) + \frac{3}{4}C_3(e_n + d_n)^2 + \frac{1}{2}C_4(e_n + d_n)^3 + \cdots} \\ &= \frac{1}{f'(x^*)} \left[ 1 - C_2(e_n + d_n) - \left( \frac{3}{4}C_3 - C_2^2 \right) (e_n + d_n)^2 + \cdots \right]. \end{aligned} \quad (8)$$

Furthermore, we have

$$f(x_n) = f'(x^*)[e_n + C_2e_n^2 + C_3e_n^3 + \cdots]. \quad (9)$$

Thus it follows (7), (8), (9) and  $y_n = x_n - v_{n-1}f(x_n)$  that

$$\begin{aligned} d_n &= x_n - x^* - v_{n-1}f(x_n) \\ &= e_n - \left[ 1 - C_2(e_{n-1} + d_{n-1}) - \left( \frac{3}{4}C_3 - C_2^2 \right) (e_{n-1} + d_{n-1})^2 + \cdots \right] [e_n + C_2e_n^2 + C_3e_n^3 + \cdots] \\ &= C_2e_n(e_{n-1} + d_{n-1}) + \left( \frac{3}{4}C_3 - C_2^2 \right) e_n(e_{n-1} + d_{n-1})^2 - C_2e_n^2 + C_2^2e_n^2(e_{n-1} + d_{n-1}) + \cdots \end{aligned} \quad (10)$$

and hence, we obtain

$$\begin{aligned} w_n &= x_n - x^* - v_nf(x_n) \\ &= e_n - \left[ 1 - C_2(e_n + d_n) - \left( \frac{3}{4}C_3 - C_2^2 \right) (e_n + d_n)^2 + \cdots \right] [e_n + C_2e_n^2 + C_3e_n^3 + \cdots] \\ &= C_2e_nd_n + \left( \frac{3}{4}C_3 - C_2^2 \right) e_n(e_n + d_n)^2 - C_2^2e_n^2(e_n + d_n) + \cdots \\ &= C_2e_nd_n + \left[ \frac{3}{4}C_3e_n + \left( \frac{3}{4}C_3 - C_2^2 \right) d_n \right] e_n(e_n + d_n) + \cdots. \end{aligned} \quad (11)$$

From (6), we obtain

$$e_{n+1} = w_n - \frac{(d_n - w_n)^2}{\alpha_1 e_{n-1} + \alpha_2 d_{n-1} + (2 - \alpha_1 - \alpha_2)w_{n-1} - \beta_1 e_n - \beta_2 d_n - (2 - \beta_1 - \beta_2)w_n}. \quad (12)$$

We first consider the case  $\alpha_1 = 0, \alpha_2 \neq 0$ , and in this case we have

$$\begin{aligned} e_{n+1} &= C_2e_nd_n - \frac{1}{\alpha_2} \frac{C_2e_n \left( \frac{e_{n-1}}{d_{n-1}} + 1 \right) + \cdots}{1 + \frac{2-\alpha_2}{\alpha_2} \frac{w_{n-1}}{d_{n-1}} + \cdots} (d_n - w_n) + \cdots \\ &= C_2e_nd_n - \frac{e_nd_n}{\alpha_2 e_{n-2}} + \cdots = -\frac{C_2e_n^2 e_{n-1}}{\alpha_2 e_{n-2}} + \cdots. \end{aligned} \quad (13)$$

From (11)–(13), we can see that, from  $z_n$  to  $x_{n+1}$ , the order is not improved when  $\alpha_1 = 0, \alpha_2 \neq 0$ . Similarly, it is obtained that the case  $\alpha_1 = \alpha_2 = 0$  also cannot improve the order. We now turn to consider the case  $\alpha \neq 0$ , and using (10)–(12), we obtain

$$\begin{aligned} e_{n+1} &= C_2e_nd_n - \frac{1}{\alpha_1} \frac{C_2e_n \left( \frac{d_{n-1}}{e_{n-1}} + 1 \right) + \cdots}{1 + \frac{\alpha_2}{\alpha_1} \frac{d_{n-1}}{e_{n-1}} + \cdots} (d_n - w_n) + \cdots \\ &= \left( 1 - \frac{1}{\alpha_1} \right) C_2e_nd_n - \frac{1}{\alpha_1} \left( 1 - \frac{\alpha_2}{\alpha_1} \right) C_2e_nd_n \frac{d_{n-1}}{e_{n-1}} + \cdots \end{aligned}$$

$$\begin{aligned}
&= \left(1 - \frac{1}{\alpha_1}\right) C_2 e_n d_n - \frac{1}{\alpha_1} \left(1 - \frac{\alpha_2}{\alpha_1}\right) C_2 e_n d_n e_{n-2} + \cdots \\
&= \left(1 - \frac{1}{\alpha_1}\right) C_2 e_n d_n + \left[1 - \frac{1}{\alpha_1} \left(2 - \frac{\alpha_2}{\alpha_1}\right)\right] C_2^3 e_n^2 e_{n-1} e_{n-2} + \cdots
\end{aligned} \tag{14}$$

From (14), we can see that the order will be improved by taking  $\alpha_1 = \alpha_2 = 1$ . In the following, by letting  $\alpha_1 = \alpha_2 = 1$ , and from (10)–(12), we obtain

$$\begin{aligned}
e_{n+1} &= w_n - \frac{d_n - w_n}{e_{n-1} + d_{n-1} - \beta_1 e_n - \beta_2 d_n - (2 - \beta_1 - \beta_2) w_n} (d_n - w_n) \\
&= C_2 e_n d_n - (d_n - w_n) \frac{C_2 e_n (e_{n-1} + d_{n-1}) + \left(\frac{3}{4} C_3 - C_2^2\right) e_n (e_{n-1} + d_{n-1})^2 + \cdots}{e_{n-1} + d_{n-1} - \beta_1 e_n - \beta_2 d_n - (2 - \beta_1 - \beta_2) w_n} + \cdots \\
&= C_2 e_n d_n - \frac{C_2 e_n + \left(\frac{3}{4} C_3 - C_2^2\right) e_n (e_{n-1} + d_{n-1}) + \cdots}{1 - \frac{\beta_1 e_n - \beta_2 d_n - (2 - \beta_1 - \beta_2) w_n}{e_{n-1} + d_{n-1}}} (d_n - w_n) + \cdots \\
&= C_2 e_n d_n - \left[ C_2 e_n + \left(\frac{3}{4} C_3 - C_2^2\right) e_n (e_{n-1} + d_{n-1}) + \cdots \right] \\
&\quad \times \left[ 1 + \frac{\beta_1 e_n - \beta_2 d_n - (2 - \beta_1 - \beta_2) w_n}{e_{n-1} + d_{n-1}} + \cdots \right] (d_n - w_n) + \cdots \\
&= C_2 e_n d_n - \left[ C_2 e_n + \left(\frac{3}{4} C_3 - C_2^2\right) e_n (e_{n-1} + d_{n-1}) + \cdots \right] \left[ 1 + \frac{\beta_1 e_n + \cdots}{e_{n-1} + d_{n-1}} + \cdots \right] (d_n - w_n) + \cdots \\
&= C_2 e_n d_n - \left[ C_2 e_n + \left(\frac{3}{4} C_3 - C_2^2\right) e_n (e_{n-1} + d_{n-1}) + \cdots \right] \left[ 1 + \frac{\beta_1 \frac{e_n}{e_{n-1}}}{1 + \frac{d_{n-1}}{e_{n-1}}} + \cdots \right] (d_n - w_n) + \cdots \\
&= C_2 e_n d_n - \left[ C_2 e_n + \left(\frac{3}{4} C_3 - C_2^2\right) e_n (e_{n-1} + d_{n-1}) + \cdots \right] \left[ 1 + \beta_1 \frac{e_n}{e_{n-1}} \left(1 - \frac{d_{n-1}}{e_{n-1}}\right) + \cdots \right] (d_n - w_n) + \cdots \\
&= C_2 e_n d_n - \left[ C_2 e_n + \left(\frac{3}{4} C_3 - C_2^2\right) e_n e_{n-1} + C_2 \beta_1 \frac{e_n^2}{e_{n-1}} + \cdots \right] (d_n - w_n) + \cdots \\
&= \left( C_2^2 - \frac{3}{4} C_3 \right) e_n e_{n-1} d_n - C_2 \beta_1 \frac{e_n^2}{e_{n-1}} d_n + \cdots \\
&= C_2 \left( C_2^2 - \frac{3}{4} C_3 \right) e_n^2 e_{n-1}^2 - C_2^2 \beta_1 e_n^3 + \cdots \\
&= C_2 \left( C_2^2 - \frac{3}{4} C_3 \right) e_n^2 e_{n-1}^2 + \cdots
\end{aligned} \tag{15}$$

Let  $A = C_2 \left( C_2^2 - \frac{3}{4} C_3 \right)$ , then (15) becomes

$$e_{n+1} = A e_n^2 e_{n-1}^2 + \cdots \tag{16}$$

Suppose that the order of (6) is  $q$  when  $\alpha_1 = \alpha_2 = 1$ , then from (4) we have

$$e_n = C e_{n-1}^q + \cdots \tag{17}$$

and

$$e_{n+1} = C e_n^q + \cdots = C^{q+1} e_{n-1}^{q^2} + \cdots \tag{18}$$

Substituting (17) and (18) into (16) gives

$$C^{q+1} e_{n-1}^{q^2} = A C^2 e_{n-1}^{2q+2} + \cdots \tag{19}$$

which implies that

$$q^2 = 2q + 2. \tag{20}$$

It is obtained from (20) that the asymptotic convergence order  $q = 1 + \sqrt{3}$ .  $\square$

**Table 1**Comparison of various iterative methods for [Example 1](#).

	Secant method	Newton's method	New method
$n$	$x_{-1} = -2, x_0 = -1$	$x_0 = -1$	$x_0 = -1$
$ x_n - x_{n-1} $	8	5	4
$ f(x_n) $	3.34199334872665e-012	2.22044604925031e-016	2.22044604925031e-016
	3.5527136788005e-015	2.66453525910038e-015	2.66453525910038e-015

**Table 2**Comparison of various iterative methods for [Example 2](#).

	Secant method	Newton's method	New method
$n$	$x_{-1} = 2, x_0 = 1.5$	$x_0 = 1.5$	$x_0 = 1.5$
$ x_n - x_{n-1} $	6	4	3
$ f(x_n) $	2.44204656496549e-012	2.22044604925031e-016	2.22044604925031e-016
	3.33066907387547e-016	4.44089209850063e-016	4.44089209850063e-016

**Table 3**Comparison of various iterative methods for [Example 3](#).

	Secant method	Newton's method	New method
$n$	$x_{-1} = 3.05, x_0 = 2.98$	$x_0 = 2.98$	$x_0 = 2.98$
$ x_n - x_{n-1} $	7	4	3
$ f(x_n) $	3.29958282918597e-013	2.22044604925031e-015	0
	0	0	0

By [Theorem 1](#), we take  $\alpha_1 = \alpha_2 = 1$  in (6), and obtain the present method given by

$$\begin{cases} y_n = x_n - v_{n-1}f(x_n), \\ v_n = f' \left( \frac{x_n + y_n}{2} \right)^{-1} \\ z_n = x_n - v_n f(x_n), \\ x_{n+1} = z_n - \frac{(y_n - z_n)^2}{x_{n-1} + y_{n-1} - \beta_1 x_n - \beta_2 y_n - (2 - \beta_1 - \beta_2)z_n} \end{cases} \quad (21)$$

where  $\beta_1, \beta_2 \in \mathbb{R}$ .

Per iteration the present methods require two evaluations of the function and one of its first derivative. We consider the definition of efficiency index [5] as  $p^{1/\omega}$ , where  $p$  is the order of the method and  $\omega$  is the number of function evaluations per iteration required by the method. We have that the present methods have the efficiency index equal to  $\sqrt[3]{1.618} \approx 1.17398$ , which is better than the method proposed in [16]  $\sqrt[5]{1.618} \approx 1.10102$ .

#### 4. Numerical example

Now we employ the new method given by (21) with  $\beta_1 = \beta_2 = 0$  and  $v_{-1} = 1$  to solve some nonlinear equations and compare them with the secant method and Newton's method defined by (1) proposed in [1–4]. The iterative method is stopped when  $|f(x)| < 1e-14$  or  $|x_n - x_{n-1}| < 1e-14$ . The examples are as follows.

##### Example 1.

$$f(x) = xe^{x^2} - \sin^2 x + 3 \cos x + 5, \quad x^* = -1.20764782713092.$$

The results of this problem are displayed in [Table 1](#).

##### Example 2.

$$f(x) = \sin^2 x - x^2 + 1, \quad x^* = 1.40449164821534.$$

The results of this problem are displayed in [Table 2](#).

##### Example 3.

$$f(x) = e^{x^2+7x-30} - 1, \quad x^* = 3.$$

The results of this problem are displayed in [Table 3](#).

From the tables, we see that the new method is efficient. It converges faster than not only the secant method but also Newton's method. In view of this fact, the new method can be viewed as a significant improvement compared with the previously known methods.

## 5. Conclusions

We present a new iterative method for solving nonlinear equations. **Theorem 1** shows that the asymptotic convergence order of this method is  $1 + \sqrt{3}$ . The numerical example shows that the method is efficient.

## Acknowledgment

The authors are grateful to the reviewer who gave the helpful suggestions.

## References

- [1] A.M. Ostrowski, *Solution of Equations and Systems of Equations*, Academic Press, New York, 1973, p. 11.
- [2] J.F. Traub, *Iterative Methods for the Solution of Equations*, Prentice-Hall, Englewood Cliffs, NJ, 1964.
- [3] J.E. Dennis, R.B. Schnable, *Numerical methods for Unconstrained Optimisation and Nonlinear Equations*, Prentice Hall, 1983.
- [4] Alfio Quarteroni, Riccardo Sacco, Fausto Saleri, *Numerical Mathematics*, Springer Science + Business Media, Inc., 2000.
- [5] W. Gautschi, *Numerical Analysis An Introduction*, Birkhäuser, Basel, 1997.
- [6] A. Melman, Geometry and convergence of Euler's and Halley's methods, *SIAM Rev.* 39 (4) (1997) 728–735.
- [7] J. Gerlach, Accelerated convergence in Newton's method, *SIAM Rev.* 36 (1994) 272–276.
- [8] S. Amat, S. Busquier, J.M. Gutiérrez, Geometric constructions of iterative functions to solve nonlinear equations, *J. Comput. Appl. Math.* 157 (2003) 197–205.
- [9] V. Kanwar, J.R. Sharma, Mamta, A new family of Secant-like method with super-linear convergence, *Appl. Math. Comput.* 171 (2005) 104–107.
- [10] H.H.H. Homeier, On Newton-type methods with cubic convergence, *J. Comput. Appl. Math.* 176 (2005) 425–432.
- [11] Kou Jisheng, Li Yitian, Wang Xiuhua, Third-order modification of Newton's method, *J. Comput. Appl. Math.* 205 (2007) 1–5.
- [12] A.Y. Özban, Some new variants of Newton's method, *Appl. Math. Lett.* 17 (2004) 677–682.
- [13] S. Weerakoon, T.G.I. Fernando, A variant of Newton's method with accelerated third-order convergence, *Appl. Math. Lett.* 13 (8) (2000) 87–93.
- [14] T.R. Scavo, J.B. Thoo, On the geometry of Halley's method, *Amer. Math. Monthly* 102 (1995) 417–426.
- [15] Hui Zhang, De-Sheng Li, Yu-Zhong Liu, A new method of secant-like for nonlinear equations, *Commun. Nonlinear Sci. Numer. Simul.* 14 (2009) 2923–2927.
- [16] Xiuhua Wang, Jisheng Kou, Chuanqing Gu, A new modified secant-like method for solving nonlinear equations, *Comput. Math. Appl.* 60 (2010) 1633–1638.
- [17] Hongmin Ren, Qingbiao Wu, Weihong Bi, On convergence of a new secant-like method for solving nonlinear equations, *Appl. Math. Comput.* 217 (2010) 583–589.
- [18] R.F. King, Tangent method for nonlinear equations, *Numer. Math.* 18 (1972) 298–304.
- [19] W. Werner, über ein Verfahren der Ordnung  $1 + \sqrt{2}$  zur Nullstellenbestimmung, *Numer. Math.* 32 (1979) 333–342.